

Purified Anti-Mouse CD106 Antibody[M/K-2.7]

catalog number: E-AB-F1091A

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


| | |
|-------------------|---|
| Reactivity | Mouse |
| Immunogen | Recombinant Mouse CD106 protein |
| Host | Rat |
| Isotype | Rat IgG1, κ |
| Clone | M/K-2.7 |
| Buffer | Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling. |

Applications

Recommended Dilution

| | |
|------------|--|
| FCM | 2 $\mu\text{g}/\text{mL}$ (0.5×10^6 - 1×10^6 cells) |
|------------|--|

Data



C57/BL6 Mouse bone marrow cells were stained with 0.2 μg Purified Anti-Mouse CD106 Antibody[M/K-2.7] (Right) and 0.2 μg Rat IgG1, κ Isotype Control (Left), followed by PE-conjugated Goat Anti-Rat IgG Secondary Antibody.

Preparation & Storage

| | |
|-----------------|--|
| Storage | Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles. |
| Shipping | Ice bag |

Background

VCAM-1 (CD106), a member of the immunoglobulin superfamily, is a cell surface protein expressed by activated endothelial cells and certain leukocytes (such as macrophages). VCAM-1 expression is induced by IL-1 beta, IL-4, TNF-alpha and IFN-gamma. VCAM-1 binds to leukocyte integrins VLA-4 and alpha 4 beta 7. The human and mouse VCAM-1 proteins share approximately 76% amino acid similarity.

For Research Use Only