Elabscience®

TRIM23 Polyclonal Antibody

catalog number: E-AB-18298

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Reactivity Human;Mouse;Rat mmnunogen Fusion protein of human TRIM23 lost Rabbit sotype IgG Purification Antigen affinity purification Conjugation Unconjugated Buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. Applications Recommended Dilution HC 1:35-1:200 Data Immunohistochemistry of paraffin-embedded Human Inmunohistochemistry of paraffin-embedded Human Immunohistochemistry of paraffin-embedded Human olorectal cancer tissue using TRIM23 Polyclonal Antibody at dilution of 1:55(×200) cervical cancer tissue using TRIM23 Polyclonal Antibody at dilution of 1:55(×200) treparation & Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. Fipping The product is shipped with ice pack,upon receipt,store it immediately at the	Note: Centinuge before opening to ensure complete recovery of viar contents.			
mmunogenFusion protein of human TRIM23lostRabbitsotypeIgGurificationAntigen affinity purificationConjugationUnconjugatedBufferPhosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.upplicationsRecommended DilutionHC1:35-1:200DataImmunohistochemistry of paraffin-embedded HumanImmunohistochemistry of paraffin-embedded HumanImmunohistochemistry of paraffin-embedded Humanolorectal cancer tissue using TRIM23 Polyclonal Antibody at dilution of 1:55(×200)Immunohistochemistry of paraffin-embedded Humanreparation & StorageStorageStorageStorageStorage / StorageStorage / Storage	Description			
Inset Rabbit sotype IgG Purification Antigen affinity purification Conjugation Unconjugated Buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. Applications Recommended Dilution HC 1:35-1:200 Pata Immunohistochemistry of paraffin-embedded Human Immunohistochemistry of paraffin-embedded Human Immunohistochemistry of paraffin-embedded Human olorectal cancer tissue using TRIM23 Polyclonal Antibody at dilution of 1:55(×200) dilution of 1:55(×200) reparation & Storage Stora at -20°C Valid for 12 months. Avoid freeze / thaw cycles. Hipping Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.	Reactivity	Human;Mouse;Rat		
sotypeIgGPurificationAntigen affinity purificationConjugationUnconjugatedBufferPhosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.ApplicationsRecommended DilutionHC1:35-1:200ConjugationImpunois for paraffinity purification solution and paraffinity purification solution and paraffinity provide solution and paraffinity provide solution and paraffinity provide solution pH 7.4, containing 0.05% stabilizer and 50% glycerol.ApplicationsRecommended DilutionHC1:35-1:200Interpret Solution provide solut	Immunogen	Fusion protein of human TRIM23		
Antigen affinity purification Conjugation Unconjugated Buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. Applications Recommended Dilution HC 1:35-1:200 Data Immunohistochemistry of paraffin-embedded Human olorectal cancer tissue using TRIM23 Polyclonal Antibody at dilution of 1:55(×200) Impunohistochemistry of paraffin-embedded Human olorectal cancer tissue using TRIM23 Polyclonal Antibody at dilution of 1:55(×200) Creparation & Storage Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. The product is shipped with ice pack,upon receipt,store it immediately at the	Host	Rabbit		
Conjugation Unconjugated Buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. Applications Recommended Dilution HC 1:35-1:200 Data Immunohistochemistry of paraffin-embedded Human Immunohistochemistry of paraffin-embedded Human Immunohistochemistry of paraffin-embedded Human olorectal cancer tissue using TRIM23 Polyclonal Antibody cervical cancer tissue using TRIM23 Polyclonal Antibody reparation & Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. Fibping The product is shipped with ice pack,upon receipt,store it immediately at the	Isotype	IgG		
Buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. Applications Recommended Dilution HC 1:35-1:200 Data Immunohistochemistry of paraffin-embedded Human Immunohistochemistry of paraffin-embedded Human Immunohistochemistry of paraffin-embedded Human olorectal cancer tissue using TRIM23 Polyclonal Antibody at dilution of 1:55(×200) Immunohistochemistry of paraffin-embedded Human reparation & Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. Fibping The product is shipped with ice pack,upon receipt,store it immediately at the	Purification	Antigen affinity purification		
Applications Recommended Dilution HC 1:35-1:200 Data Immunohistochemistry of paraffin-embedded Human Immunohistochemistry of paraffin-embedded Human Immunohistochemistry of paraffin-embedded Human Ioorectal cancer tissue using TRIM23 Polyclonal Antibody at dilution of 1:55(×200) Immunohistochemistry of paraffin-embedded Human Preparation & Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. Fibiping The product is shipped with ice pack,upon receipt,store it immediately at the	Conjugation	Unconjugated		
HC 1:35-1:200 Data Immunohistochemistry of paraffin-embedded Human olorectal cancer tissue using TRIM23 Polyclonal Antibody at dilution of 1:55(×200) dilution of 1:55(×200) Preparation & Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. Shipping The product is shipped with ice pack,upon receipt,store it immediately at the	Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.		
Data Immunohistochemistry of paraffin-embedded Human olorectal cancer tissue using TRIM23 Polyclonal Antibody at dilution of 1:55(×200) Cervaration & Storage Store at -20°C Valid for 12 months. A void freeze / thaw cycles. The product is shipped with ice pack,upon receipt,store it immediately at the	Applications	Recommended Dilution		
Immunohistochemistry of paraffin-embedded Human Immunohistochemistry of paraffin-embedded Human olorectal cancer tissue using TRIM23 Polyclonal Antibody at dilution of 1:55(×200) Immunohistochemistry of paraffin-embedded Human Preparation & Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. Shipping Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.	IHC	1:35-1:200		
olorectal cancer tissue using TRIM23 Polyclonal Antibody cervical cancer tissue using TRIM23 Polyclonal Antibody at dilution of 1:55(×200) dilution of 1:55(×200) Preparation & Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. The product is shipped with ice pack,upon receipt,store it immediately at the	Data			
StorageStore at -20°C Valid for 12 months. Avoid freeze / thaw cycles.ShippingThe product is shipped with ice pack,upon receipt,store it immediately at the	colorectal cancer tissue using TRIM23 Polyclonal Antibody cervical cancer tissue using TRIM23 Polyclonal Antibody a			
hipping The product is shipped with ice pack,upon receipt,store it immediately at the	Preparation & Storage			
	Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.		
termerature recommended	Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the		
temperature recommended.				

Background

The protein encoded by this gene is a member of the tripartite motif (TRIM) family. The TRIM motif includes three zincbinding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. This protein is also a member of the ADP ribosylation factor family of guanine nucleotide-binding family of proteins. Its carboxy terminus contains an ADP-ribosylation factor domain and a guanine nucleotide binding site, while the amino terminus contains a GTPase activating protein domain which acts on the guanine nucleotide binding site. The protein localizes to lysosomes and the Golgi apparatus. It plays a role in the formation of intracellular transport vesicles, their movement from one compartment to another, and phopholipase D activation. Three alternatively spliced transcript variants for this gene have been described.