

## Recombinant Human PD-1/PDCD1 Protein (His Tag)

Catalog Number: PKSH031642

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	HEK293 Cells-derived Human PD-1/PDCD1 protein Met 1-Gln 167, with an C-terminal His
<b>Calculated MW</b>	17.4 kDa
<b>Accession</b>	NP_005009.2
<b>Bio-activity</b>	Immobilized human PD-1 at 10 µg/ml (100 µl/well) can bind recombinant human B7-H1 / PD-L1 / Fc chimera with a linear range of 0.02-0.4 µg/ml.


### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



Lot-to-Lot Consistency




Immobilized human PD-1 (Cat: PKSH031642) at 2 µg/ml (100 µl/well) can bind PD-L1 Protein, Human, Recombinant (ECD, hFc Tag) (Cat: PKSH031887), the EC<sub>50</sub> of human B7-H1 is 150-600 ng/mL.

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017



Captured Anti-PD-1 Mab (Human IgG4) on proA Chip can bind Recombinant Human PD-1 Protein, His Tag (Cat: PKSH031642) with an affinity constant of 10.03 nM as determined in a SPR assay (Biacore T200) .

## Background

Programmed cell death 1; also known as PDCD1; is a type I transmembrane glycoprotein; and is an immunoreceptor belonging to the CD28/CTLA-4 family negatively regulates antigen receptor signaling by recruiting protein tyrosine phosphatase; SHP-2 upon interacting with either of two ligands; PD-L1 or PD-L2. PD1 inhibits the T-cell proliferation and production of related cytokines including IL-1; IL-4; IL-10 and IFN- $\gamma$ ; by suppressing the activation and transduction of PI3K/AKT pathway. In addition; coligation of PD1 inhibits BCR-mediated signal by dephosphorylating key signal transducer. PD1 has been suggested to be involved in lymphocyte clonal selection and peripheral tolerance; and thus contributes to the prevention of autoimmune diseases. Furthermore; PD1 is shown to be a regulator of virus-specific CD8+ T cell survival in HIV infection. As a cell surface molecule; PDCD1 regulates the adaptive immune response. Engagement of PD-1 by its ligands PD-L1 or PD-L2 transduces a signal that inhibits T-cell proliferation; cytokine production; and cytolytic function.

## For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017