

Recombinant Human IL-18 protein (His tag)

Catalog Number:PDEH100217



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	IGIF;IL-18;IL-1g;IL1F4;Interleukin 18;IL18;IGIF
Species	Human
Expression Host	E.coli
Sequence	Tyr 37-Asp 193
Accession	Q14116
Calculated Molecular Weight	17.2 kDa
Observed molecular weight	18 kDa
Tag	C-His

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01 % Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Interleukin-18 (IL-18; also known as interferon-gamma inducing factor) is a proinflammatory cytokine that belongs to the IL-1 superfamily and is produced by macrophages and other cells. This cytokine can induce the IFN-gamma production of T cells. The combination of IL-18 and IL12 has been shown to inhibit IL4 dependent IgE and IgG1 production; and enhance IgG2a production of B cells. IL-18 binding protein (IL18BP) can specifically interact with this cytokine; and thus negatively regulate its biological activity. IL-18 is an IL-1-like cytokine that requires cleavage with caspase-1 to become active; was found to increase IgE production in a CD4+ T cells-; IL-4- and STAT6-dependent fashion. IL-18 and T cell receptor-mediated stimulation could induce naïve CD4+ T cells to develop into IL-4-producing cells in vitro. Thus; caspase-1 and IL-18 may be critical in regulation of IgE production in vivo; providing a potential therapeutic target for

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Human IL-18 protein (His tag)

Catalog Number:PDEH100217



allergic disorders. IL-18 production in primary synovial cultures and purified synovial fibroblasts was; in turn; upregulated by TNF- α and IL-1 β ; suggesting that monokine expression can feed back to promote Th1 cell development in synovial membrane. Besides; synergistic combinations of IL-18; IL-12; and IL-15 may be of importance in sustaining both Th1 responses and monokine production in RA.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com