Elabscience®

IGHA1 Polyclonal Antibody


catalog number: E-AB-52649


Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description		
Reactivity	Human	
Immunogen	Fusion protein of human IGHA1	
Host	Rabbit	
Isotype	IgG	
Purification	Antigen affinity purification	
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.	

Applications	Recommended Dilution
IHC	1:50-1:300

Data





Immunohistochemistry of paraffin-embedded Human cervicalImmunohistochemistry of paraffin-embedded Humancancer tissue using IGHA1 Polyclonal Antibody at dilution of colorectal cancer tissue using IGHA1 Polyclonal Antibody at $1:50(\times 200)$ dilution of $1:50(\times 200)$ dilution of $1:50(\times 200)$

Preparation & Storage	
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

Constant region of immunoglobulin heavy chains. Immunoglobulins, also known as antibodies, are membrane-bound or secreted glycoproteins produced by B lymphocytes. In the recognition phase of humoral immunity, the membranebound immunoglobulins serve as receptors which, upon binding of a specific antigen, trigger the clonal expansion and differentiation of B lymphocytes into immunoglobulins-secreting plasma cells. Secreted immunoglobulins mediate the effector phase of humoral immunity, which results in the elimination of bound antigens (PubMed:22158414, PubMed: 20176268). The antigen binding site is formed by the variable domain of one heavy chain, together with that of its associated light chain. Thus, each immunoglobulin has two antigen binding sites with remarkable affinity for a particular antigen. The variable domains are assembled by a process called V-(D)-J rearrangement and can then be subjected to somatic hypermutations which, after exposure to antigen and selection, allow affinity maturation for a particular antigen (PubMed:17576170, PubMed:20176268). Ig alpha is the major immunoglobulin class in body secretions (PubMed:2241915)

For Research Use Only