

Purified Anti-Human CD277 Antibody[ICT01]

catalog number: AN008880P

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description


Reactivity	Human
Immunogen	Recombinant Human CD277 protein
Host	Mouse
Isotype	Human IgG1, κ
Clone	ICT01
Purification	>98%, Protein A/G purified
Buffer	Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling.

Applications

Recommended Dilution

FCM	2 µg/mL (0.5×10 ⁶ -1×10 ⁶ cells)
------------	--

Data



HEK293T cells transfected with pcDNA3.1 plasmid encoding human CD277 gene were stained with 0.2 µg Purified Anti-Human CD277 Antibody[ICT01] (Right) and 0.2 µg Human IgG1, κ Isotype Control (Left), followed by APC-conjugated Goat Anti-Human IgG Secondary Antibody.

Preparation & Storage

Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
Shipping	Ice bag

Background

Butyrophilin subfamily 3A (BTN3A), also known as CD277, is a 56 kD type 1 transmembrane glycoprotein. The butyrophilin (BT)/B7 3A family includes three members, BTN3A1, BTN3A2, and BTN3A3, which share a highly conserved extracellular IgV-like domain. BTN3A is broadly expressed in various immune cells including T cells, B cells, NK cells, monocytes, dendritic cells, endothelial cells, and some tumor cells. It plays a role in T cell activation, regulating their proliferation and release of IFN and other cytokines.

For Research Use Only