# Elabscience®

### Monkey PLAU/uPA Antibody Pair Set

Catalog No.	E-KAB-0660	Applications
Synonyms	UPA;u-PA;ATF;URK;UP-A;Urokinase;Abbokinase	

### ELISA

#### **Kit components & Storage**

Title	Specifications	Storage
Monkey PLAU/uPA Capture Antibody	1 vial, 100 µ g	Store at -20°C for one year. Avoid
		freeze/thaw cycles.
Monkey PLAU/uPA Detection Antibody	1 vial, 50 μL	Store at -20°C for one year. Avoid
(Biotin)		freeze/thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Product Information**

Items		Characteristic (E-KAB-0660)	
		Monkey PLAU/uPA Capture	Monkey PLAU/uPA Detection
		Antibody	Antibody (Biotin)
Immunogen	Immunogen	Recombinant Monkey PLAU/uPA	Recombinant Monkey PLAU/uPA
Information		protein	protein
	Swissprot	Q9GK78	
Product details	Reactivity	Monkey	Monkey
	Host	Mouse	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5 mg/mL	/
	Buffer	PBS with 0.04% Proclin 300; 50%	PBS with 0.04% Proclin 300; 1%
		glycerol; pH 7.5	protective protein; 50% glycerol; pH
			7.5
	Purify	Protein A or G	Antigen Affinity
	Specificity	Detects Monkey PLAU/uPA in ELISAs.	

For Research Use Only

## **Elabscience**®

#### Applications

Monkey PLAU/uPA Sandwich ELISA Assay

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4 μg/mL	Monkey PLAU/uPA Capture	
Capture		Antibody	10
			Optical Density
ELISA	1:1000-1:10000	Monkey PLAU/uPA	Optic
Detection		Detection Antibody (Biotin)	
			0. 1 1 1000 10000 1000 Monkey PLAU/uPA Concentration(pg/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

#### Background

This gene encodes a serine protease involved in degradation of the extracellular matrix and possibly tumor cell migration and proliferation. A specific polymorphism in this gene may be associated with late-onset Alzheimer's disease and also with decreased affinity for fibrin-binding. This protein converts plasminogen to plasmin by specific cleavage of an Arg-Val bond in plasminogen. Plasmin in turn cleaves this protein at a Lys-Ile bond to form a two-chain derivative in which a single disulfide bond connects the amino-terminal A-chain to the catalytically active , carboxy-terminal B-chain. This two-chain derivative is also called HMW-uPA (high molecular weight uPA) . HMW-uPA can be further processed into LMW-uPA (low molecular weight uPA) by cleavage of chain A into a short chain A (A1) and an amino-terminal fragment. LMW-uPA is proteolytically active but does not bind to the uPA receptor. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

For Research Use Only