

Recombinant Mouse IL1R1/CD121a Protein (His Tag)

Catalog Number: PKSM041327

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Mouse
Source	HEK293 Cells-derived Mouse IL1R1/CD121a protein Leu20-Lys338, with an C-terminal His
Calculated MW	38.1 kDa
Observed MW	50-90 kDa
Accession	P13504
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
Reconstitution	Please refer to the specific buffer information in the printed manual. Please refer to the printed manual for detailed information.

Data



Background

For Research Use Only

Mouse Interleukin 1 receptor, type I (IL-1R1) also known as CD121a (Cluster of Differentiation 121a), is an interleukin receptor. IL-1R1/CD121a is a cytokine receptor that belongs to the interleukin 1 receptor family. This protein is a receptor for interleukin 1 alpha (IL1A), interleukin 1 beta (IL1B), and interleukin 1 receptor antagonist (IL1RA). It is an important mediator involved in many cytokine induced immune and inflammatory responses. An IL1 receptor accessory protein that can heterodimerize with the Type I receptor in the presence of IL1 α or IL1 β but not IL1ra, was identified. This Type I receptor complex appears to mediate all the known IL1 biological responses. The receptor Type II has a short cytoplasmic domain and does not transduce IL1 signals. In addition to the membranebound form of IL1 RII, a naturally occurring soluble form of IL1 RII has been described. It has been suggested that the Type II receptor, either as the membranebound or as the soluble form, serves as a decoy for IL1 and inhibits IL1 action by blocking the binding of IL1 to the signaling Type I receptor complex.