## **Elabscience**®

### **OPRK1** Polyclonal Antibody

### catalog number: E-AB-64902


Note: Centrifuge before opening to ensure complete recovery of vial contents.


1:50-1:200

Description	
Reactivity	Human;Mouse;Rat
Immunogen	A synthetic peptide of human OPRK1 (NP_000903.2).
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution
WB	1:500-1:2000

#### Data


IHC






OPRK1 Polyclonal Antibody at dilution of 1:1000.

**Observed-MW:40 kDa** Calculated-MW:33 kDa/42 kDa



Western blot analysis of extracts of various cell lines using Immunohistochemistry of paraffin-embedded Rat testis using OPRK1 Polyclonal Antibody at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded Human stomach using OPRK1 Polyclonal Antibody at dilution of 1:100 (40x lens).

Immunohistochemistry of paraffin-embedded Mouse kidney using OPRK1 Polyclonal Antibody at dilution of 1:100 (40x lens).

Preparation & Storage	
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

For Research Use Only

Toll-free: 1-888-852-8623 Web:www.elabscience.com

Tel: 1-832-243-6086 Email:techsupport@elabscience.com Fax: 1-832-243-6017

# **Elabscience**®

This gene encodes an opioid receptor, which is a member of the 7 transmembrane-spanning G protein-coupled receptor family. It functions as a receptor for endogenous ligands, as well as a receptor for various synthetic opioids. Ligand binding results in inhibition of adenylate cyclase activity and neurotransmitter release. This opioid receptor plays a role in the perception of pain and mediating the hypolocomotor, analgesic and aversive actions of synthetic opioids. Variations in this gene have also been associated with alcohol dependence and opiate addiction. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. A recent study provided evidence for translational readthrough in this gene and expression of an additional C-terminally extended isoform via the use of an alternative in-frame translation termination codon.

For Research Use Only