Elabscience®

UBTF Polyclonal Antibody

catalog number: E-AB-92999

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description			
Reactivity	Human;Mouse;Rat	Human;Mouse;Rat	
Immunogen	Recombinant fusion prot	Recombinant fusion protein of human UBTF	
Host	Rabbit	Rabbit	
Isotype	IgG	IgG	
Purification	Affinity purification	Affinity purification	
Buffer	Phosphate buffered solut	tion, pH 7.4, containing 0.05% stabilizer and 50% glycerol.	
Applications	Recommended Dilu	Recommended Dilution	
WB	1:500-1:2000	1:500-1:2000	
Data			
UBTF Polyclonal	extracts of various cell lines using Antibody at 1:1000 dilution.	^{170kDa} ^{100kDa} ^{100kDa} ^{70kDa} ^{40kDa} ^{40kDa} ^{40kDa} ^{80kDa} ^{80kDa} ^{100kDa ^{100kDa} ^{100kDa} ^{100kDa} ^{100kDa} ^{100kDa} ^{100kDa} ^{100kDa} ^{100kDa} ^{100kDa} ^{100kDa} ^{100kDa} ^{100kDa} ^{100kDa} ^{100kDa} ^{100kDa} ^{100kDa} ^{100kDa} ^{100kDa} ^{100kDa} }	
Observed-MW:84 kDa		Observed-MW:84 kDa	
Calculated-MW:84 kDa/89 kDa		Calculated-MW:84 kDa/89 kDa	
Preparation & Storage			
Storage	Store at -20°C Valid for 12	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.	
Shipping	The product is shipped v	The product is shipped with ice pack, upon receipt, store it immediately at the	
	temperature recommended.		
Rookground			

Background

This gene encodes a member of the HMG-box DNA-binding protein family. The encoded protein plays a critical role in ribosomal RNA transcription as a key component of the pre-initiation complex, mediating the recruitment of RNA polymerase I to rDNA promoter regions. The encoded protein may also play important roles in chromatin remodeling and pre-rRNA processing, and its activity is regulated by both phosphorylation and acetylation. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. Pseudogenes of this gene are located on the short arm of chromosomes 3, 11 and X and the long arm of chromosome 11.