ARIH2 Polyclonal Antibody

catalog number: E-AB-52978



Note: Centrifuge before opening to ensure complete recovery of vial contents.

5	
Description	
Reactivity	Human;Mouse
Immunogen	Fusion protein of human ARIH2
Host	Rabbit
Isotype	IgG
Purification	Antigen affinity purification
Conjugation	Unconjugated
buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution
IHC	1:50-1:300
Data	
tissue using ARIH2 Pc	f paraffin-embedded Human tonsil olyclonal Antibody at dilution of
	1:70(×200)
Preparation & Storage	
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.
Background	
TRIAD1, also known as finger and two RING-typ TRIAD1 interacts with U from E2 ubiquitin-conjug subject to post-translatio TRIAD1 maps to human	ARIH2 (ariadne homolog 2) or ARI2, is a 493 amino acid protein that contains one IBR-type zinc pe zinc fingers and belongs to the ariadne subfamily of RBR proteins. Localized to the nucleus, JBE2L3 and is thought to act as an E3 ubiquitin-protein ligase, functioning to accept ubiquitin gating enzymes and transfer the acquired ubiquitin residue to target substrates. TRIAD1 is onal DNA damage-dependent phosphorylation, probably by ATM or ATR. The gene encoding chromosome 3, which houses over 1,100 genes, including a chemokine receptor (CKR) gene numan cancer-related gene loci.

For Research Use Only