

Recombinant Human Fetuin-A/AHSG Protein (His Tag)

Catalog Number: PDMH100405

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*


Description

Species	Human
Source	HEK293 Cells-derived Human Fetuin-A protein Met1-Val367, with an C-terminal His
Calculated MW	40.3 kDa
Observed MW	58 kDa
Accession	P02765
Bio-activity	Not validated for activity

Properties

Purity	> 95% as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU/mg of the protein as determined by the LAL method
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Human Fetuin-A/AHSG proteins, 2 µg/lane of Recombinant Human Fetuin-A/AHSG proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 58 kDa.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Fetuin-A, also known as Alpha-2-HS-Glycoprotein (AHSG), belongs to the Fetuin family, is a plasma binding protein, and is more abundant in fetal than adult blood. It is involved in several functions, such as endocytosis, brain development and the formation of bone tissue. Fetuins are carrier proteins like albumin. Fetuin-A forms soluble complexes with calcium and phosphate and thus is a carrier of otherwise insoluble calcium phosphate. Thus Fetuin-A is a potent inhibitor of pathological calcification. The circulating levels of fetuin-A, a well-described inhibitor of calcification, regulate the cell-dependent process of osteogenesis. The low circulating fetuin-A levels are associated with a greater prevalence and/or severity of Vascular calcification (VC) and increased risk for all-cause and cardiovascular mortality. However, high circulating fetuin-A levels appear to induce insulin resistance and, in non-dialyzed subjects with diabetic nephropathy, are directly related to VC burden. The emerging role of fetuin-A deficiency as a risk factor in dialysis patients was documented in cross-sectional studies demonstrating a significant correlation with all-cause and cardiovascular mortality. Additionally, Human fetuin-A is a negative acute phase protein involved in inflammatory diseases, thus being a potential physiological regulator of meprin activity. Fetuin-A is a broad-range protease inhibitor. Fetuin-A and cystatin C as endogenous proteolytic regulators of meprin activity broadens our understanding of the proteolytic network in plasma.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017