Recombinant Human/Mouse/Rat Irisin/FNDC5 (N-6His)

Catalog Number: PKSM041407



Description			
Species	Human/Mouse/Rat		
Mol_Mass	13.4 kDa		
Accession	Q8NAU1		
Bio-activity	Not validated for activity		
Properties			
Purity	>95% as determined by reducing SDS-PAGE.		
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.		
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80		
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of		
	reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.		
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.		
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.		
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants		
	before lyophilization.		
	Please refer to the specific buffer information in the printed manual.		
Reconstitution	Please refer to the printed manual for detailed information.		

Note: Centrifuge before opening to ensure complete recovery of vial contents.

kDa	MK	R
120 90 60		
40	-	•
30	-	-
20	-	U
14	_	-

Background

Data

Fibronectin type III domain-containing protein 5, the precursor of irisin, is a protein that is encoded by the FNDC5 gene. Human Irisin is synthesized as a 212 amino acid (aa) precursor encoding a type 1 transmembrane protein with a 121 aa extracellular domain (ECD), a 21 aa transmembrane domain, and a 39 aa cytoplasmic domain. The ECD of Irisin contains a fibronectin type III domain and multiple glycosylation sites. The ECD is proteolytically cleaved to release the 112 aa soluble Irisin hormone into circulation.Mature human, mouse share 100% sequence identity.Irisin induces expression of peroxisome proliferatoractivated receptor γ coactivator 1 α (PGC1 α) and uncoupling protein1(UCP1), mitochondrialassociated metabolic proteins. Irisin induces the transition of white adipose tissue into more metabolically

mitochondrialassociated metabolic proteins. Irisin induces the transition of white adipose tissue into more metabolically active beige adipose tissue. Irisin also regulates neuronal cell differentiation and neurite outgrowth in the brain and is involved in the differentiation of osteoblasts.

For Research Use Only