

A Reliable Research Partner in Life Science and Medicine

# APC Anti-Human CD11b Antibody[HI11b]

Catalog Number: E-AB-F1317E

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

Reactivity Human Mouse Host

Isotype Mouse IgG2b, ĸ

Clone No. HI11b

APC Mouse IgG2b, κ Isotype Control[MPC-11] [Product E-AB-F09812E] Isotype Control

APC Conjugation

**Conjugation Information** APC is designed to be excited by the Red (627-640 nm) laser and detected using an

optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein

protectant.

## **Applications**


#### Recommended usage

**FCM** 

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

#### individual use.

#### **Data**



Staining of normal human peripheral blood cells with APC Anti-Human CD11b Antibody[HI11b] (filled gray histogram) or APC Mouse IgG2b, κ Isotype Control (empty black histogram). Cells in the granulocytes gate were used for analysis.

## **Preparation & Storage**

**Storage** Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

**Shipping** Ice bag

## **Antigen Information**

**Uniprot ID** P11215 Gene ID 3684

# For Research Use Only

# Elabscience®

#### **Elabscience Bionovation Inc.**

A Reliable Research Partner in Life Science and Medicine

## **Background**

CD11b is a 165-170 kD type I transmembrane glycoprotein also known as  $\alpha$ Mintegrin, Mac-1, CR3, and C3biR. CD11b non-covalently associates with integrin  $\beta$ 2(CD18) and is expressed on granulocytes, monocytes/macrophages, dendritic cells, NK cells, and subsets of T and B cells. CD11b/CD18 is critical for the transendothelial migration of monocytes and neutrophils. It is also involved in granulocyte adhesion, phagocytosis, and neutrophil activation. CD11b/CD18 interacts with ICAM-1 (CD54), ICAM-2 (CD102), ICAM-4, CD14, CD23, heparin, iC3b, fibrinogen, and factor X.

Fax: 1-832-243-6017