

Recombinant Human Fcγ RIIA/CD32a Protein(Fc Tag)

Catalog Number: PDMH100334

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	HEK293 Cells-derived Human Fcγ RIIA/CD32a proteins Gln34-Gly217, with an C-terminal Fc
Calculated MW	45.1 kDa
Observed MW	50 kDa
Accession	P12318
Bio-activity	Not validated for activity

Properties

Purity	> 90% as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU/mg of the protein as determined by the LAL method
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Human Fcγ RIIA/CD32a proteins, 2 µg/lane of Recombinant Human Fcγ RIIA/CD32a proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 50 KD

Background

For Research Use Only

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Human Fc γ Rs are divided into three classes designated Fc γ RI (CD64); Fc γ RII (CD32); and Fc γ RIII (CD16); which generate multiple isoforms; are recognized. The activating- type receptor either has or associates non-covalently with an accessory subunit that has an immunoreceptor tyrosine-based activation motif (ITAM) in its cytoplasmic domain. Fc γ RI binds IgG with high affinity and functions during early immune responses; whereas Fc γ RII and RIII are low affinity receptors that recognize IgG as aggregates surrounding multivalent antigens during late immune responses. Human CD32; also known as Low affinity immunoglobulin γ Fc region receptor II-a (IgG Fc receptor II-a); Fc γ RII A or FCGR2A Protein; is expressed on cells of both myeloid and lymphoid lineages as well as on cells of non-hematopoietic origin. Associated with an ITAM-bearing adapter subunit; FcR γ ; CD32a (Fc γ RII A) delivers an activating signal upon ligand binding; and results in the initiation of inflammatory responses including cytosis; phagocytosis; degranulation; and cytokine production. The responses can be modulated by signals from the co-expressed inhibitory receptors such as Fc γ RII B; and the strength of the signal is dependent on the ratio of expression of the activating and inhibitory receptors.

For Research Use Only

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017