

PNOC Polyclonal Antibody

catalog number: E-AB-16651

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human
Immunogen	Synthetic peptide of human PNOC
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.


Applications Recommended Dilution

IHC	1:50-1:200
------------	------------

Data



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using PNOC Polyclonal Antibody at dilution 1:40



Immunohistochemistry of paraffin-embedded Human brain tissue using PNOC Polyclonal Antibody at dilution 1:40

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

Nociception, a pain response mechanism, occurs in response to stimuli that threaten the integrity of an organism. The first synapses produced as a result of the initiation of nociception are modulated by excitatory amino acids (glutamate and aspartate) and many peptides (substance P, CGRP, CCK, endogenous opioids). Nociceptin (also designated orphanin FQ) is a neuronal peptide that is similar to opioid peptides. Nociceptin activates KOR-3 (kappa-type opioid receptor, also designated ORL1), a Gprotein-coupled receptor. Although similar to dynorphin A, a kappa opioid peptide, nociceptin functions to make animals hyperreactive to nociceptive stimulations. Nociceptin is also involved in locomotor behavior and may be involved in the modulation of synaptic plasticity in learning and memory.

For Research Use Only