Recombinant Mouse TROP2/TACSTD2 Protein (His Tag)

Catalog Number: PKSM041262

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description		
Species	Mouse	
Source	HEK293 Cells-derived Mouse TROP2/TACSTD2 protein Gln25-Gly270, with an C-	
	terminal His	
Calculated MW	28.8 kDa	
Observed MW	38-55 kDa	
Accession	Q8BGV3	
Bio-activity	Not validated for activity	
Properties		
Purity	> 95 % as determined by reducing SDS-PAGE.	
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.	
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80	
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of	
	reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.	
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.	
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.	
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants	
	before lyophilization.	
	Please refer to the specific buffer information in the printed manual.	
Reconstitution	Please refer to the printed manual for detailed information.	
Data		

kDa	MK	R
120 90 60 40		
30	-	
20)	
14	-	

Background

Tumor-associated calcium signal transducer 2(Tacstd2), also known as Cell surface glycoprotein Trop-2, belongs to the EPCAM family. Tacstd2 expressed in kidney, lung, ovary and testis and has high levels of expression in immortalized keratinocytes. Tacstd2 may functions as a growth factor receptor. It has negative regulation of branching involved in ureteric bud morphogenesis, cell motility, epithelial cell migration, ruffle assembly, stress fiber assembly and substrate adhesion-dependent cell spreading. Also, it has positive regulation of stem cell proliferation. Tacstd2 is capable of transducing an intracellular calcium signal and may play a role in tumor growth. It also has adhesive functions.

For Research Use Only Toll-free: 1-888-852-8623