

## PE/Cyanine7 Anti-Human CD55 Antibody[HI55a]

Catalog Number: AN00336H

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description

<b>Reactivity</b>	Human
<b>Host</b>	Mouse
<b>Isotype</b>	Mouse IgG2a, κ
<b>Clone No.</b>	HI55a
<b>Isotype Control</b>	PE/Cyanine7 Mouse IgG2a, κ Isotype Control[C1.18.4] [Product E-AB-F09802H]
<b>Conjugation</b>	PE/Cyanine 7
<b>Conjugation Information</b>	PE/Cyanine7 is designed to be excited by the Blue (488 nm), Green (532 nm) and yellow-green (561 nm) lasers and detected using an optical filter centered near 775 nm (e.g., a 780/60 nm bandpass filter).
<b>Storage Buffer</b>	Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.

### Applications

#### Recommended usage

<b>FCM</b>	Each lot of this antibody is quality control tested by flow cytometric analysis. <b>The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood).</b> Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.
------------	---

### Preparation & Storage

<b>Storage</b>	Keep as concentrated solution. This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.
<b>Shipping</b>	Ice bag

### Antigen Information

<b>Alternate Names</b>	Complement decay-accelerating factor;CD55;CR;DAF
<b>Uniprot ID</b>	P13591
<b>Gene ID</b>	1604
<b>Background</b>	CD55 is a 60-70 kD glycosylphosphatidylinositol (GPI)-anchored single chain glycoprotein also known as decay-accelerating factor (DAF). It is expressed on hematopoietic cells including erythrocytes and many non-hematopoietic cells. CD55 accelerates the dissociation of the components of the C3-convertases (namely C2a from C4b in the C4bC2a complex, a C3-convertase of the classical pathway, and factor Bb from the C3bBb complex, a C3-convertase of the alternative pathway) to protect cells from inappropriate damage caused by autologous complement. CD55 has been reported to reduce the efficiency of NK cell lysis and induce signal transduction in T cells. CD55 has also been shown to interact with CD97 and bind to Coxsackie and Echovirus.

### For Research Use Only