

IL-4/BSF-1, Human, Recombinant

Cat. No. : PCK031

产品信息

别名	BCGF;BCDF;B-cell Stimulating Factor (BSF-1)
物种	Human
表达宿主	E.coli
序列信息	MHKCDITLQEIIKTLNSLTEQKTLCTELTVTDIFAASKNTTEKETFCRAATVLRQFYSHHEKDTR CLGATAQQFHRHKQLIRFLKRLDRNLWGLAGLNSCPVKEANQSTLENFLERLKTIMREKYSKC SS with polyhistidine tag at the C-terminus.
检索号	P05112.1
标签	His-tag at the C-terminus
分子量	15.9 kDa
有效期	12 months
生物活性	Measure by its ability to induce TF-1 cells proliferation. The ED50 for this effect is < 0.2 ng/mL. The specific activity of recombinant human IL-4 is approximately > 2.8 × 10 ⁷ IU/mg.

产品特性


内毒素 (EU/μg)	< 0.1
保存	Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10% FBS, 5% HSA or 5% trehalose solution), protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.
运输	Ambient temperature or ice pack.
制剂	The protein was lyophilized from a 0.2 μm filtered solution containing 1 × PBS, pH 8.0.
复溶	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 μg/mL. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

背景介绍

The interleukin 4 (IL4, IL-4) is a cytokine that induces differentiation of naive helper T cells (Th0 cells) to Th2 cells. Upon activation by IL-4, Th2 cells subsequently produce additional IL-4 in a positive feedback loop. The cell that initially produces IL-4, thus inducing Th2 differentiation, has not been identified, but recent studies suggest that basophils may be the effector cell. It is closely related and has functions similar to Interleukin 13.



SDS-PAGE



普诺赛® | Procell system

普诺赛® | Procell system

普诺赛® | Procell system

