

**TGFB1/TGFβ1/TGF-β1, Human, Recombinant**

Cat. No. : PCK091

**产品信息**

别名	Transforming Growth Factor Beta-1;TGF-Beta-1;Latency-Associated Peptide;LAP;TGFB1;TGFB
物种	Human
表达宿主	CHO Stable Cells
序列信息	Ala279-Ser390
检索号	P01137
分子量	12.8 kDa
有效期	12 months
生物活性	Measured by its ability to inhibit the IL-4-dependent proliferation of TE-1 human erythroleukemic cells. The ED50 for this effect is 4-40 pg/mL.

**产品特性**


内毒素	< 1.0 EU per 1 µg as determined by LAL test.
保存	Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -5~-20°C for 3 months.
运输	Ambient temperature or ice pack.
制剂	Lyophilized from a 0.2 µm filtered solution of 50 mM Glycine-HCl, 150 mM NaCl, 4% Mannitol, pH 2.5.
复溶	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/mL. Dissolve the lyophilized protein in sterile water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**背景介绍**


Transforming Growth Factor β-1 (TGFβ-1) is a secreted protein which belongs to the TGF-β family. TGFβ-1 is abundantly expressed in bone, articular cartilage and chondrocytes and is increased in osteoarthritis (OA). TGFβ-1 performs many cellular functions, including the control of cell growth, cell proliferation, cell differentiation and apoptosis. The precursor is cleaved into a latency-associated peptide (LAP) and a mature TGFβ-1 peptide. TGFβ-1 may also form heterodimers with other TGFβ family members. It has been found that TGFβ-1 is frequently upregulated in tumor cells. Mutations in this gene results in Camurati-Engelmann disease.



## SDS-PAGE



## Bio activity



Measured by its ability to inhibit the IL-4-dependent proliferation of TF-1 human erythroleukemic cells. The ED50 for this effect is 4-40 pg/mL.

