



A Reliable Research Partner in Life Science and Medicine

Mouse IL-3 Antibody Pair Set

Catalog No. E-KAB-0695 Applications ELISA

Synonyms IL3;MCGF;MCSF;MULTI-CSF;P-Cell Stimulating Factor

Kit components & Storage

Title	Specifications	Storage
Mouse IL-3 Capture Antibody	1 vial, 100 μ g	Store at -20°C. Avoid freeze / thaw
		cycles.
Mouse IL-3 Detection Antibody (Biotin)	1 vial, 50 μL	Store at -20°C. Avoid freeze / thaw
		cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0695)	
		Mouse IL-3 Capture Antibody	Mouse IL-3 Detection Antibody
			(Biotin)
Immunogen	Immunogen	Recombinant Mouse IL-3 protein	Recombinant Mouse IL-3 protein
Information	Swissprot	P01586	
Product details	Reactivity	Mouse	Mouse
	Host	Goat	Rat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5mg/mL	/
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300, 1%
		glycerol, pH 7.4	protective protein, 50% glycerol, pH
			7.4
	Purify	Affinity purification	Affinity purification
	Specificity	Detects Mouse IL-3 in ELISAs.	

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com Email: techsupport@elabscience.com

Fax: 1-832-243-6017



A Reliable Research Partner in Life Science and Medicine

Applications

Mouse IL-3 Sandwich ELISA Assay

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4ug/mL	Mouse IL-3 Capture Antibody	
Capture			10
			» 13
			Optical Density
ELISA	1:1000-1:10000	Mouse IL-3 Detection	ond O 0.1
Detection		Antibody (Biotin)	· ·
			0.01
			Mouse IL-3 concentration (pg/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

Interleukin-3 (IL-3) is a hemopoietic growth factor involved in the survival, proliferation, and differentiation of multipotent hemopoietic cells. In addition to granulocyte-macrophage colony-stimulating factor (GMCSF) and IL-5, IL-3 is the most potent growth factor for basophils. On mature basophils, these three cytokines act through specific receptors mediating adhesion, migration, and releasability. Mast cells expresses high amounts of IL-3. In the absence of antigen, rapid release of large amount of IL-3 in an autocrine dependent manner is responsible for mast cell survival. IL-3 plays a vital role in stimulating basophils and mast cell responses to parasite infections. IL-3 has also been implicated in the pathogenesis of several chronic inflammatory diseases, including asthma, atherosclerosis, and neurodegenerative disorders, such as multiple sclerosis. IL-3 stimulates colony formation of megakaryocytes, neutrophils, and macrophages from bone marrow cultures. IL-3 is expressed in the major embryonic vessels and regulates the survival and proliferation of hematopoietic stem cells in the early stages of embryonic development.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com