

A Reliable Research Partner in Life Science and Medicine

Elab Fluor® Red 780 Anti-Mouse CD19 Antibody[1D3]

Catalog Number: E-AB-F0986S

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse Host Rat

lsotype Rat lgG2a, κ

Clone No. 1D3

Isotype Control Elab Fluor® Red 780 Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09832S]

Conjugation Elab Fluor[®] Red 780

Conjugation Information Elab Fluor[®] Red 780 is designed to be excited by the Red (627-640 nm) laser and

detected using an optical filter centered near 770 nm (e.g., a 780/60 nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein

protectant.

Applications


Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

individual use

Data



Mouse splenocytes are stained with Elab Fluor[®] Red 780 Anti-Mouse CD19 Antibody (filled gray histogram). Unstained splenocytes (blank black histogram) are used as control.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

Rev. V1.6

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names B-lymphocyte antigen CD19;CD19;Cd19;Differentiation antigen CD19

 Uniprot ID
 P25918

 Gene ID
 12478

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web:www.elabscience.com
 Email:techsupport@elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Background

CD19 is a 95 kD glycoprotein also known as B4. It is a member of the lg superfamily, expressed on all pro-B to mature B cells (during development) and follicular dendritic cells. Plasma cells do not express CD19. CD19, in association with CD21 and CD81, forms a molecular complex integral to B cell activation.

Fax: 1-832-243-6017