## **Elabscience**®

### Recombinant Human CCL21 Protein(Trx Tag)

### Catalog Number: PDEH100535

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description			
Species	Human		
Source	E.coli-derived Human CCL21 protein Ser24-Pro134, with an N-terminal Trx		
Calculated MW	32 kDa		
Observed MW	35 kDa		
Accession	O00585		
Bio-activity	Not validated for activity		
Properties			
Purity	> 90% as determined by reducing SDS-PAGE.		
Endotoxin	< 10 EU/mg of the protein as determined by the LAL method		
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80		
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at $<$ -20°C for 3 months.		
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.		
Formulation	Lyophilized from a 0.2 $\mu$ m filtered solution in PBS with 5% Trehalose and 5%		
	Mannitol.		
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of		
	0.5 mg/mL. Concentration is measured by UV-Vis.		

Data

М	R
-	
_	-
	M

SDS-PAGE analysis of Human CCL21 proteins, 2  $\mu$ g/lane of Recombinant Human CCL21 proteins was resolved with an SDS-PAGE under reducing conditions, showing bands at 32

#### KD

Background

# **Elabscience**®

This antimicrobial gene is one of several CC cytokine genes clustered on the p-arm of chromosome 9. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. Similar to other chemokines the protein encoded by this gene inhibits hemopoiesis and stimulates chemotaxis. This protein is chemotactic in vitro for thymocytes and activated T cells, but not for B cells, macrophages, or neutrophils. The cytokine encoded by this gene may also play a role in mediating homing of lymphocytes to secondary lymphoid organs. It is a high affinity functional ligand for chemokine receptor 7 that is expressed on T and B lymphocytes and a known receptor for another member of the cytokine family (small inducible cytokine A19).