PE/Cyanine7 Anti-Human CD18 Antibody[TS1/18.1.2.11]

Catalog Number: E-AB-F1057H

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human
Host	Mouse
Isotype	Mouse IgG1, κ
Clone No.	TS1/18.1.2.11
Isotype Control	PE/Cyanine7 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792H]
Conjugation	PE/Cyanine 7
Conjugation Information Storage Buffer	PE/Cyanine7 is designed to be excited by the Blue (488 nm), Green (532 nm) and yellow-green (561 nm) lasers and detected using an optical filter centered near 775 nm (e.g., a 780/60 nm bandpass filter). Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.
Applications	Recommended usage
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Preparation & Storage	
Storage	Keep as concentrated solution.
	This product can be stored at 2-8 $^\circ$ C for 12 months. Please protected from prolonged
	exposure to light and do not freeze.
Shipping	Ice bag
Antigen Information	
Alternate Names	CD18;Cell surface adhesion glycoproteins LFA-1/CR3/p150+95 subunit beta;
	Complement receptor C3 subunit beta;Integrin beta-2;Itgb2
Uniprot ID	P05107
Gene ID	3689
Background	CD18 is a 90-95 kD type I transmembrane protein also known as integrin $\beta 2$ subunit,
	LFA-1 β subunit, and β 2 integrin. CD18 non-covalently associates with CD11a, CD11b
	or CD11c. CD18 is expressed on all leukocytes. CD18 and associated α chains
	function in adhesion and signaling in hematopoietic cells.

For Research Use Only