

A Reliable Research Partner in Life Science and Medicine

Purified Anti-Human CD14 Antibody[H332-1B10]

catalog number: AN002130P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human

Immunogen Recombinant Human sCD14 protein expressed by Mammalian

Host Rat

Isotype Rat IgG1, ĸ Clone H332-1B10

Purification >98%, Protein A/G purified

Buffer Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer.


Dialyze to completely remove the stabilizer prior to labeling.


Recommended Dilution Applications

 $0.1-0.4 \mu g/mL$ **ELISA Detector**

FCM $2 \mu g/mL(0.5 \times 10^6 - 1 \times 10^6 \text{ cells})$

Data





Human peripheral blood monocytes were stained with 0.2 µg 0.2 μg Rat IgG1, κ Isotype Control (Left), followed by PEconjugated Goat Anti-Rat IgG Secondary Antibody.

Sandwich ELISA-Recombinant Human sCD14 protein Purified Anti-Human CD14 Antibody[H332-1B10] (Right) and standard curve.Background subtracted standard curve using sCD14 antibody(AN002120P)(Capture),sCD14 antibody(AN002130P)(Detector) in sandwich ELISA.The reference range value for Recombinant Human sCD14 protein is 1.25-80 ng/mL.

Rev. V2.1

Preparation & Storage

Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

For Research Use Only

Toll-free: 1-888-852-8623 Fax: 1-832-243-6017 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com

Elabscience®

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

The cluster of differentiation (CD) system is commonly used as cell markers in immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules associating with the immune function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. Cluster of differentiation 14 (CD14) is a member of the CD system. It takes its name from its inclusion in the CD molecule surface marker proteins. CD14 exists in two forms: a form anchored into the membrane or a soluble form. CD14 was found expressed in macrophages, neutrophil granulocyte and dendritic cells. The major function is to serve as a co-receptor (along with TLR4 and MD-2) for the bacterial lipopolysaccharide (LPS) and other pathogenassociated molecular patterns.

For Research Use Only