

Recombinant Human STING/TMEM173 Protein (Sumo & His Tag)



Catalog Number:PKSH033517

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Stimulator of interferon genes protein;TMEM173;Mediator of IRF3 activation;sting;
Species	Human
Expression Host	E.coli
Sequence	Val155-Val341
Accession	Q86WV6
Calculated Molecular Weight	33.8 kDa
Observed molecular weight	35 kDa
Tag	N-SUMO-His

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
Shipping	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < - 20°C.
Formulation	Supplied as a 0.2 µm filtered solution of 20mM HEPES, 100mM NaCl, 10% Glycerol, pH8.0.
Reconstitution	Not Applicable

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Stimulator of Interferon Gene(Sting,TMEM173) belongs to the TMEM173 family. STING is 379 amino acids (aa) in length. It contains an N-terminal cytoplasmic region (aa 1-20), four transmembrane segments (aa 21-173), and a C-terminal cytoplasmic domain (aa 174-379). It ubiquitously expressed in skin endothelial cells, alveolar type 2 pneumocytes, bronchial epithelium and alveolar macrophagesand. Its subunit structure associated with the MHC-II complex and Interacts with DDX58/RIG-I, MAVS and SSR2, RNF5 and TRIM56 along with TBK1. This type of protein often uses as facilitator of innate immune signaling that acts as a sensor of cytosolic DNA from bacteria and viruses and promotes the production of type I interferon.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017