

Recombinant Phospho-eIF2α (Ser51) Monoclonal Antibody

catalog number: AN300145L

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description



Reactivity	Human;Mouse;Rat
Immunogen	A synthetic phosphopeptide corresponding to residues around Ser51 of human Phospho-eIF2α.
Host	Rabbit
Isotype	IgG
Clone	A1259
Purification	Protein A
Buffer	10 mM sodium HEPES, 150 mM NaCl, 100 µg/mL protein protectant, 50% glycerol, pH 7.5

Applications

Recommended Dilution


IHC	1:100-1:1000
WB	1:2000-1:10000
IF	1:200-1:1000
ELISA	1:5000-1:20000

Data



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-eIF2α (Phospho Ser51) antibody. The HRP-conjugated Goat anti-Rabbit IgG (H + L) antibody was used to detect the antibody. Lane 1: HeLa Lane 2: HeLa was treated with Thapsigargin (300nM) for 30 minutes Predicted band size: 36kDa Observed band size: 36kDa

Human hepatocellular carcinoma was stained with anti-eIF2α (Phospho Ser51) Rabbit antibody



Immunofluorescence analysis of A549. 1, primary Antibody (red) was diluted at 1:200 (4°C overnight). 2, Goat Anti Rabbit IgG (H&L) - Alexa Fluor 594 Secondary antibody was diluted at 1:1000 (room temperature, 50min). 3, DAPI (blue) 10min.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.4

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	Ice bag

Background

The translation initiation factor EIF2 catalyzes the first regulated step of protein synthesis initiation, promoting the binding of the initiator tRNA to 40S ribosomal subunits. Binding occurs as a ternary complex of methionyl-tRNA, EIF 2, and GTP. EIF2 is composed of 3 nonidentical subunits, the 36-kD EIF2-alpha subunit (EIF2S1), the 38-kD EIF2-beta subunit (EIF2S2; MIM 603908), and the 52-kD EIF2-gamma subunit (EIF2S3; MIM 300161). The rate of formation of the ternary complex is modulated by the phosphorylation state of EIF2-alpha (Ernst et al., 1987 [PubMed 2948954]).[supplied by OMIM, Feb 2010]

For Research Use Only