

Recombinant Mouse Contactin 2/CNTN2 Protein (His Tag)

Catalog Number: PKSM040349

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Mouse
Source	HEK293 Cells-derived Mouse Contactin 2/CNTN2 protein Gln31-Glu 1013, with an C-terminal His
Calculated MW	108.7 kDa
Observed MW	110 kDa
Accession	Q61330
Bio-activity	Not validated for activity

Properties

Purity	> 97 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 97 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Contactins are a subgroup of molecules belonging to the immunoglobulin superfamily that are expressed exclusively in the nervous system. The subgroup consists of six members: Contactin-1, Contactin-2(TAG-1), Contactin-3(BIG-1), BIG-2, Contactin-5(NB-2) and NB-3. Axonal expression and the neurite extension activity of Contactin-1 and Contactin-2 attracted researchers to study the function of these molecules in axon guidance during development. Contactin-1 and Contactin-2 have come to be known as the principal molecules in the function and maintenance of myelinated neurons. Contactin-2, also known as CNTN2, is a glycosylphosphatidylinositol (GPI)-anchored neuronal membrane protein that functions as a cell adhesion molecule. Contactin-2 is expressed by a subset of neuronal populations in the developing central nervous system (CNS) and peripheral nervous system (PNS). Contactin-2 is also expressed by oligodendrocytes and Schwann cells, which are myelinating glial cells of the CNS and PNS, respectively. Contactin-2 may play a role in the formation of axon connections in the developing nervous system. Contactin-2 is also involved in glial tumorigenesis and may provide a potential target for therapeutic intervention. During embryonic development, Contactin-2 interacts either in a homophilic, or heterophilic fashion with various transmembrane proteins.