PAICS Polyclonal Antibody

catalog number: E-AB-15282



Note: Centrifuge before opening to ensure complete recovery of vial contents.

				7.0	
	06	TO T	m	11	on
v				ш	w

Reactivity Human; Mouse; Rat

Immunogen Recombinant protein of human PAICS

Host Rabbit
Isotype IgG

PurificationAffinity purificationConjugationUnconjugated

buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.


Ap	plications	Recommended Dilution

WB 1:500-1:2000 **IHC** 1:50-1:200

Data


KDa 1 2 3 4 130— 95— 72— 55— 36—

Western Blot analysis of Hela, K562, Raji and Jurkat cell using PAICS Polyclonal Antibody at dilution of 1:550



Immunohistochemistry of paraffin-embedded Human cervical cancer using PAICS Polyclonal Antibody at dilution of 1:50

Calculated-MV:47 kDa



Immunohistochemistry of paraffin-embedded Human esophagus cancer using PAICS Polyclonal Antibody at dilution of 1:50

Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

This gene encodes a bifunctional enzyme containing phosphoribosylaminoimidazole carboxylase activity in its N-terminal region and phosphoribosylaminoimidazole succinocarboxamide synthetase in its C-terminal region. It catalyzes steps 6 and 7 of purine biosynthesis. The gene is closely linked and divergently transcribed with a locus that encodes an enzyme in the same pathway, and transcription of the two genes is coordinately regulated. The human genome contains several pseudogenes of this gene. Multiple transcript variants encoding different isoforms have been found for this gene.

For Research Use Only