

## ZAP70 Polyclonal Antibody

catalog number: E-AB-53219

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

|                     |  |
|---------------------|--|
| <b>Reactivity</b>   | Human;Mouse  |
| <b>Immunogen</b>    | Synthetic peptide of human ZAP70   |
| <b>Host</b>         | Rabbit   |
| <b>Isotype</b>      | IgG  |
| <b>Purification</b> | Antigen affinity purification  |
| <b>Buffer</b>       | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |

### Applications

| Applications | Recommended Dilution |
|--------------|----------------------|
| <b>WB</b>    | 1:500-1:2000         |
| <b>IHC</b>   | 1:25-1:100           |

### Data




Western blot analysis of Jurkat cells using ZAP70 Polyclonal

Antibody at dilution of 1:1000

**Observed-MW:Refer to figures**

**Calculated-MW:70 kDa**



Immunohistochemistry of paraffin-embedded Human thyroid

cancer tissue using ZAP70 Polyclonal Antibody at dilution of

1:50(×200)

### Preparation & Storage

|                 |  |
|-----------------|--|
| <b>Storage</b>  | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.  |
| <b>Shipping</b> | The product is shipped with ice pack,upon receipt,store it immediately at the temperature recommended. |

### Background

Zeta-associated protein (ZAP)-70,a member of the Syk family of tyrosine kinases,plays an important role in T-cell receptor signaling,natural killer (NK) cell activation and early B-cell development. ZAP-70 is expressed in a subset of cases of chronic lymphocytic leukemia/small lymphocytic lymphoma (CLL/SLL) with unmutated immunoglobulin heavy-chain variable region (IgVH) genes and is associated with poor clinical outcome.

### For Research Use Only