A Reliable Research Partner in Life Science and Medicine

Recombinant Human CD3 ε/CD3E Protein (His Tag)

Catalog Number: PKSH033388

Note: Centrifuge before opening to ensure complete recovery of vial contents.

		. •
I P S	crin	non
Des	CIIP	LIUII

Species Human

Source HEK293 Cells-derived Human CD3 & CD3E protein Asp23-Asp126, with an C-terminal

His

 Mol_Mass
 12.8 kDa

 Accession
 P07766

Bio-activity Immobilized Human CD3E-His at 10 ug/ml (100 ul/wel)can bind Human Anti-CD3

mAb. The ED50 of Recombinant Human Anti-CD3 mAb is 3-15 ng/ml.

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

T-Cell Surface Glycoprotein CD3 ϵ Chain (CD3 ϵ) is a single-pass type I membrane protein. CD3 ϵ contains 1 Ig-like (immunoglobulin-like) domain and 1 ITAM domain. CD3 ϵ is a polypeptide encoded by the CD3E gene on chromosome 11 in humans. The T cell receptor-CD3 complex (TCR/CD3 complex) is involved in T-cell development and several intracellular signal-transduction pathways. This complex is critical for T-cell development and function, and represents one of the most complex transmembrane receptors. The T cell receptor-CD3 complex is unique in having ten cytoplasmic immunoreceptor tyrosine-based activation motifs (ITAMs). TCR/CD3 complex plays an important role in coupling antigen recognition to several intracellular signal-transduction pathways.

For Research Use Only