

Elab Fluor® 647 Anti-Rat CD38 Antibody[14.27]

Catalog Number: AN00959M

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Rat
Host	Mouse
Isotype	Mouse IgG2b, κ
Clone No.	14.27
Isotype Control	Elab Fluor® 647 Mouse IgG2b, κ Isotype Control[MPC-11] [Product E-AB-F09812M]
Conjugation	Elab Fluor® 647
Conjugation Information	Elab Fluor® 647 is designed to be excited by the Red laser (627-640 nm) and detected using an optical filter centered near 670 nm (e.g., a 660/20 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications

Recommended usage

FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 µL staining volume or per 100 µL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.
-----	---

Data



Staining of Rat splenocytes cells with FITC Anti-Rat CD3

Antibody and Elab Fluor® 647 Anti-Rat CD38
Anyibody[14.27] (left) or Elab Fluor® 647 Mouse IgG2b, κ
Isotype Control (right). Total viable cells were used for
analysis.

Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 12 months. Please protect from prolonged exposure to light and do not freeze.
Shipping	Ice bag

Antigen Information

Alternate Names	T10;ADP-ribosyl cyclase
Uniprot ID	Q64244
Gene ID	25668

For Research Use Only

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.3

Background

CD38 is a 45 kD type II transmembrane ADP-ribosyl cyclase expressed on early hematopoietic precursors and leukocytes, hepatic stellate cells, astrocytes and epithelial cells. CD38 expression can be upregulated upon cell activation. CD38 catalyzes the production of cyclic ADP ribose, which is important in Ca²⁺ release. CD38 induces production of proinflammatory cytokines and proliferation. The ligand for CD38 is CD31. Impairment in CD38 expression is associated with immunological and behavioral disorders.

For Research Use Only

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.3