

A Reliable Research Partner in Life Science and Medicine

# Elab Fluor® 700 Anti-Mouse CD4 Antibody [RM4-5]

Catalog Number: E-AB-F1353M1

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Description**

Reactivity Mouse Host Rat

Isotype Rat IgG2a, κ
Clone No. RM4-5

Isotype Control Elab Fluor® 700 Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09832M1]

Conjugation Elab Fluor® 700

**Conjugation Information** Elab Fluor® 700 is designed to be excited by the Red laser (627-640 nm) and detected

using an optical filter centered near 719 nm (e.g., a 725/40 nm bandpass filter).


Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.

#### Applications Recommended usage

**FCM** 

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

#### Data



Staining of C57BL/6 murine splenocytes with FITC Anti-

Mouse CD3 Antibody[17A2] and Elab Fluor<sup>®</sup> 700 Anti-Mouse CD4 Antibody[RM4-5](left) or Elab Fluor<sup>®</sup> 700 Rat IgG2a, κ Isotype Control(right). Total viable cells were used for analysis.

#### **Preparation & Storage**

**Storage** Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

#### **Antigen Information**

Alternate Names L3T4;T-cell surface antigen T4/Leu-3;T-cell surface glycoprotein CD4;T4

Web: www.elabscience.cn

 Uniprot ID
 P06332

 Gene ID
 12504

### For Research Use Only



## Elabscience Biotechnology Co., Ltd.

A Reliable Research Partner in Life Science and Medicine

#### **Background**

CD4 is a 55 kD protein also known as L3T4 or T4. It is a member of the Ig superfamily, primarily expressed on most thymocytes and a subset of T cells, and weakly on macrophages and dendritic cells. It acts as a co-receptor with the TCR during T cell activation and thymic differentiation by binding MHC class II and associating with the protein tyrosine kinase Ick.