

Recombinant Human LGALS7 Protein(Trx Tag)

Catalog Number: PDEH100555

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	E.coli-derived Human LGALS7 protein Ser2-Phe136, with an N-terminal Trx
Calculated MW	34.8 kDa
Observed MW	34.8 kDa
Accession	P47929
Bio-activity	Not validated for activity

Properties

Purity	> 90% as determined by reducing SDS-PAGE.
Endotoxin	< 10 EU/mg of the protein as determined by the LAL method
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Human LGALS7 proteins, 2 µg/lane of Recombinant Human LGALS7 proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 34.8 KD

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.7

LGALS7, also known as Galectin-7, is a member of the galectins family. The galectins are a family of beta-galactoside-binding proteins. There are at least 14 identified members of this family. Galectins share similarities in the CRD (the carbohydrate recognition domain). They are synthesized as cytosolic proteins. Though localized principally in the cytoplasm and lacking a classical signal peptide, galectins can also be stimulated to secretion by non-classical pathways or targeted to the nucleus. Galectins are implicated in modulating cell-cell and cell-matrix interactions. LGALS7 contains 1 galectin domain and is mainly expressed in stratified squamous epithelium. Galectin-7 could be involved in cell-cell and/or cell-matrix interactions necessary for normal growth control. LGALS7 is a pro-apoptotic protein that functions intracellularly upstream of JNK activation and cytochrome c release.