

Recombinant Human PARP-1 Protein (His Tag)

Catalog Number:PKSH031294



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	ADPRT;ADPRT1;ARTD1;pADPRT-1;PARP;PARP-1;PPOL
Species	Human
Expression Host	Baculovirus-Insect Cells
Sequence	Met 1-Trp 1014
Accession	NP_001609.2
Calculated Molecular Weight	114.5 kDa
Observed molecular weight	100-110 kDa
Tag	C-His
Bioactivity	Immobilized human PARP1 at 10 µg/mL (100 µl/well) can bind biotinylated human HSP70. The EC50 of biotinylated human HSP70 is 0.035 µg/mL.

Properties

Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

Poly (ADP-ribose) polymerase 1(PRAP1), also known as NAD(+) ADP-ribosyltransferase 1(ADPRT), is a chromatin-associated enzyme which modifies various nuclear proteins by poly(ADP-ribosylation). The ADP-D-ribosyl group of NAD⁺ is transferred to an acceptor carboxyl group on a histone or the enzyme itself, and further ADP-ribosyl groups are transferred to the 2'-position of the terminal adenosine moiety, building up a polymer with an average chain length of 20-30 units. The poly(ADP-ribosylation) modification is critical for a wide range of processes, including DNA repair, regulation of chromosome structure, transcriptional regulation, mitosis and apoptosis. PARP1 is demonstrated to mediate

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Human PARP-1 Protein (His Tag)

Catalog Number:PKSH031294



the poly(ADP-ribose) ation of APLF (aprataxin PNK-like factor) and CHFR (checkpoint protein with FHA and RING domains), two representative proteins involved in the DNA damage response and checkpoint regulation. Further, It has been suggested that DNA-dependent protein kinase (DNA-PK), another component of DNA repair, suppresses PARP activity, probably through direct binding and/or sequestration of DNA-ends which serve as an important stimulator for both enzymes. PARP1 inhibitors is thus proposed as a targeted cancer therapy for recombination deficient cancers, such as BRCA2 tumors.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017