

A Reliable Research Partner in Life Science and Medicine

PerCP/Cyanine5.5 Anti-Human CD25 Antibody[BC96]

Catalog Number: E-AB-F1194J

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human Host Mouse

Isotype Mouse IgG1, κ

Clone No. BC96

Isotype Control PerCP/Cyanine5.5 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792J]

Conjugation PerCP/Cyanine 5.5

Conjugation Information PerCP/Cyanine5.5 is designed to be excited by the blue laser (488 nm) and detected

using an optical filter centered near 675 nm (e.g., a 690/50 nm bandpass filter).


Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Human peripheral blood lymphocytes are stained with FITC Anti-Human CD4 Antibody and PerCP/Cyanine5.5 Anti-Human CD25 Antibody (Left). Lymphocytes are stained with FITC Anti-Human CD4 Antibody and PerCP/Cy5.5 Mouse IgG1, κ Isotype Control (Right).

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names IL-2 receptor subunit alpha;IL-2-RA;IL-2R subunit alpha;IL2-RA;IL2RA;Interleukin-2

receptor subunit alpha;TAC antigen;p55

Uniprot ID P01589

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web:www.elabscience.com Email:techsupport@elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Gene ID Background 3559

CD25 is a 55 kD type I transmembrane glycoprotein also known as the low affinity IL-2 receptor α chain or Tac. It is expressed on progenitor lymphocytes, activated T and B cells, and activated monocytes/macrophages. CD25 is also expressed on a subset of non-stimulated CD4+ T cells termed T regulatory cells. CD25 associates with the IL-2 receptor β (CD122) and common γ chains (CD132) to form the high affinity IL-2R complex.

Fax: 1-832-243-6017