

A Reliable Research Partner in Life Science and Medicine

Recombinant CD13/ANPEP Monoclonal Antibody

catalog number: AN300044P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human

Immunogen Recombinant Human CD13 / ANPEP protein


Host Rabbit
Isotype IgG
Clone A1154
Purification Protein A

Buffer 0.2 µm filtered solution in PBS


Applications Recommended Dilution

WB 1:500-1:2000 **IHC-P** 1:100-1:500


Data



Immunohistochemistry of paraffin-embedded human renal carcinoma using CD13 / ANPEP Monoclonal Antibody at dilution of 1:200.



Immunohistochemistry of paraffin-embedded human liver using CD13 / ANPEP Monoclonal Antibody at dilution of 1:200.



Western Blot with CD13 / ANPEP Monoclonal Antibody at dilution of 1:500. Lane A: THP1 Whole Cell Lysate, Lane B: U937 Whole Cell Lysate, Lysates/proteins at 30 µg per lane.

Observed-MW:140 kDa Calculated-MW:110 kDa

Preparation & Storage

Storage This antibody can be stored at 2°C-8°C for one month without detectable loss of

activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.

Shipping Ice bag

Background

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: w w w .elabscience.com
 Email: techsupport@elabscience.com

Elabscience®

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Aminopeptidase N is located in the small-intestinal and renal microvillar membrane, and also in other plasma membranes. In the small intestine aminopeptidase N plays a role in the final digestion of peptides generated from hydrolysis of proteins by gastric and pancreatic proteases. Its function in proximal tubular epithelial cells and other cell types is less clear. The large extracellular carboxyterminal domain contains a pentapeptide consensus sequence characteristic of members of the zinc-binding metalloproteinase superfamily. Sequence comparisons with known enzymes of this class showed that CD13 and aminopeptidase N are identical. The latter enzyme was thought to be involved in the metabolism of regulatory peptides by diverse cell types, including small intestinal and renal tubular epithelial cells, macrophages, granulocytes, and synaptic membranes from the CNS. Human aminopeptidase N is a receptor for one strain of human coronavirus that is an important cause of upper respiratory tract infections. Defects in this gene appear to be a cause of various types of leukemia or lymphoma.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com
 Rev. V1.2