

## PE Anti-Human CD204 Antibody[7C9C20]

**Catalog Number:** AN00876D

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description


<b>Reactivity</b>	Human
<b>Host</b>	Mouse
<b>Isotype</b>	Mouse IgG2a, κ
<b>Clone No.</b>	7C9C20
<b>Isotype Control</b>	PE Mouse IgG2a, κ Isotype Control[C1.18.4] [Product E-AB-F09802D]
<b>Conjugation</b>	PE
<b>Conjugation Information</b>	PE is designed to be excited by the Blue (488 nm), Green (532 nm) and Yellow-Green (561 nm) lasers and detected using an optical filter centered near 575 nm (e.g., a 585/42 nm bandpass filter).
<b>Storage Buffer</b>	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

### Applications

### Recommended usage

<b>FCM</b>	Each lot of this antibody is quality control tested by flow cytometric analysis. <b>The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood).</b> Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.
------------	---

### Data



GM-CSF-stimulated (6 days) human peripheral blood mononuclear cells surface stained with PE Anti-Human CD204 Antibody[7C9C20](left) or PE Mouse IgG1, κ Isotype Control(right).

### Preparation & Storage

<b>Storage</b>	Keep as concentrated solution. This product can be stored at 2-8°C for 12 months. Please protect from prolonged exposure to light and do not freeze.
<b>Shipping</b>	Ice bag

### Antigen Information

<b>Alternate Names</b>	Macrophage scavenger receptor;MSR;MSR1;SRA,CD204
<b>Uniprot ID</b>	P21757
<b>Gene ID</b>	4481

### For Research Use Only

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

Rev. V1.3

## Background

CD204, also known as scavenger receptor A (SR-A) and the macrophage scavenger receptor (MSR), is one of the phagocytic pattern-recognition receptors (PRRs) expressed on macrophages and dendritic cells. CD204 was initially identified as a receptor mediating recognition and internalization of low-density lipoprotein (LDL) by macrophages and playing critical roles in atherogenesis. CD204 recognizes apoptotic cells, modified lipid proteins, and exogenous pathogen-associated molecular patterns (PAMPs), which results in the induction of innate immune and inflammatory responses. CD204 can act as a co-receptor for Toll-like receptors, such as TLR3, TLR4, or TLR 9, to facilitate the expression of proinflammatory cytokines. CD204 has been implicated in several pathological processes such as Alzheimer's disease, sepsis, ischemic injury, and coronary artery disease.

## For Research Use Only

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

Rev. V1.3