

CXCL10/IP-10/Crg-2, Mouse, Recombinant**Cat. No. : GPCK335****产品信息**

物种	Mouse
表达宿主	E.coli
序列信息	IPLARTVRCNCIHIDDGPVRMRAIGKLEIPASLSCPRVEIIATMKKNDEQRCLNPESKTIKNLMK AFSQKRSKRAP with polyhistidine tag at the N-terminus.
检索号	Q548V9
标签	His-tag at the N-terminus
分子量	9.47 kDa
有效期	12 months
生物活性	Measure by its ability to chemoattract BaF3 cells transfected with human CXCR3. The ED50 for this effect is < 0.2 µg/mL.

产品特性

内毒素 (EU/µg)	< 0.1
保存	Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10% FBS, 5% HSA or 5% trehalose solution), protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.
运输	Ambient temperature or ice pack.
制剂	The protein was lyophilized from a 0.2 µm filtered solution containing 1 × PBS, pH 7.4.
复溶	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 µg/mL. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

背景介绍

CXCL10, belongs to the CXC chemokine family, is also known as IP-10. CXCL10 was originally identified as a responder to IFN-gamma in monocytes, endothelial cells and fibroblasts. The effects of this chemokine can be induced by binding to the cell surface chemokine receptor CXCR3. Research shows that CXCL10 is a chemoattractant for activated T-lymphocytes, monocytes and macrophages. It also has other functions, such as promotion of T cell adhesion to endothelial cell.

