

Recombinant CD155/PVR Monoclonal Antibody

catalog number: AN300499P

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*


Description

Reactivity	Mouse
Immunogen	Recombinant Mouse CD155/PVR protein
Host	Rabbit
Isotype	IgG
Clone	8C3
Purification	Protein A
Buffer	0.2 µm filtered solution in PBS


Applications Recommended Dilution

ICC/IF	1:50-1:1000
FCM	1:25-1:100

Data



Immunofluorescence analysis of Mouse PVR(m CD155) in raw264.7 cells. Cells were fixed with 4% PFA, blocked with 10% serum, and incubated with Rabbit anti-Mouse PVR (m CD155) monoclonal antibody (1:100) at 4°C overnight. Then cells were stained with the Alexa Fluor® 488-conjugated (left panel, captured by laser confocal scanning microscope; right panel, captured by fluorescence microscope) Goat Anti-rabbit IgG secondary antibody, countstained with DAPI (blue). Positive staining was localized to plasma membrane.



Flow cytometric analysis of Mouse PVR(CD155) expression on BABL/c splenocytes. Cells were stained with purified anti-Mouse PVR(CD155), then a FITC-conjugated second step antibody. The fluorescence histograms were derived from gated events with the forward and side light-scatter characteristics of intact cells.

Preparation & Storage

Storage	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
Shipping	Ice bag

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.0

CD155, commonly known as PVR (poliovirus receptor) and Necl-5 (nectin-like molecule-5), is a type I transmembrane single-span glycoprotein, and belongs to the nectins and nectin-like (Necl) subfamily. CD155 was originally identified based on its ability to mediate the cell attachment and entry of poliovirus (PV), an etiologic agent of the central nervous system disease poliomyelitis. The normal cellular function is in the establishment of intercellular adherens junctions between epithelial cells. CD155 may assist in an efficient humoral immune response generated within the intestinal immune system. It has been demonstrated that CD155 can be recognized and bond by DNAM-1 and CD96 which promote the adhesion, migration and NK-cell killing, and thus efficiently prime cell-mediated tumor-specific immunity.