Purified Anti-Mouse CD11a Antibody[FD441.8]

catalog number: E-AB-F1033A



Note: Centrifuge before opening to ensure complete recovery of vial contents.

-					
	AC	cri	m	Т	٦n
\mathbf{L}		CII		2.	

Reactivity Mouse Host Rat


 $\begin{array}{lll} \textbf{Isotype} & & \text{Rat IgG2b, } \kappa \\ \textbf{Clone} & & \text{FD441.8} \\ \textbf{Conjugation} & & \text{Unconjugated} \\ \end{array}$

buffer PBS, pH 7.2. Contains 0.05% proclin 300.

Applications Recommended Dilution

FCM $2 \mu g/mL(1 \times 10^5 - 5 \times 10^5 \text{ cells})$

Data



C57/BL6 Mouse splenocytes were stained with 0.2 μg Purified Anti-Mouse CD11a Antibody[FD441.8] (Right) and 0.2 μg Rat IgG2b, κ Isotype Control (Left), followed by PEconjugated Goat Anti-Rat IgG Secondary Antibody.

Preparation & Storage

Storage Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles.

Shipping Ice bag

Background

CD11a is a 180 kD glycoprotein, also known as α L integrin, LFA-1 α , Ly-15, or Ly-21. It is a member of the integrin family, primarily expressed on lymphocytes, monocytes/macrophages, and granulocytes. In association with CD18, the CD11a/CD18 complex forms LFA-1. CD11a plays an important role in intercellular adhesion and costimulation by binding its ligands, ICAM-1 (CD54), ICAM-2 (CD102), and ICAM-3 (CD50).

For Research Use Only