

## AURKAIP1 Polyclonal Antibody

catalog number: E-AB-10976

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description

<b>Reactivity</b>	Human
<b>Immunogen</b>	Recombinant protein of human AURKAIP1
<b>Host</b>	Rabbit
<b>Isotype</b>	IgG
<b>Purification</b>	Affinity purification
<b>Conjugation</b>	Unconjugated
<b>Buffer</b>	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

### Applications

### Recommended Dilution

<b>WB</b>	1:500-1:2000
-----------	--------------

### Data



Western Blot analysis of Human testis tissue using  
AURKAIP1 Polyclonal Antibody at dilution of 1:600

**Calculated-MW:22 kDa**

### Preparation & Storage

<b>Storage</b>	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
<b>Shipping</b>	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

### Background

AKIP (AURKA-interacting protein), also known as AURKAIP1 (aurora kinase A interacting protein 1) or AIP, is a 199 amino acid protein that localizes to the nucleus and is ubiquitously expressed, with highest levels present in testis, heart and skeletal muscle. Interacting specifically with ARK-1 (aurora kinase 1), AKIP functions to induce the proteasomal-dependent degradation of ARK-1, thereby acting as a negative regulator of ARK-1 activity. AKIP is encoded by a gene which maps to human chromosome 1, which spans 260 million base pairs, contains over 3,000 genes and comprises nearly 8% of the human genome. Chromosome 1 houses a large number of disease-associated genes, including those that are involved in familial adenomatous polyposis, Stickler syndrome, Parkinson's disease, Gaucher disease, schizophrenia and Usher syndrome. Aberrations in chromosome 1 are found in a variety of cancers, including head and neck cancer, malignant melanoma and multiple myeloma.

### For Research Use Only