

## Recombinant Mouse NKG2-D type II Integral Membrane Protein/NKG2D/CD314 (N-Fc)

Catalog Number: PKSM041404

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

|                      |  |
|----------------------|--|
| <b>Species</b>       | Mouse  |
| <b>Source</b>        | HEK293 Cells-derived Mouse NKG2D/CD314 protein Phe94-Val232, with an N-terminal Fc |
| <b>Calculated MW</b> | 42.2 kDa   |
| <b>Observed MW</b>   | 45-60 kDa  |
| <b>Accession</b>     | O54709   |
| <b>Bio-activity</b>  | Not validated for activity   |

### Properties

|                       |  |
|-----------------------|--|
| <b>Purity</b>         | > 95 % as determined by reducing SDS-PAGE.   |
| <b>Endotoxin</b>      | < 1.0 EU per µg of the protein as determined by the LAL method.  |
| <b>Storage</b>        | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.            |
| <b>Shipping</b>       | This product is provided as lyophilized powder which is shipped with ice packs.  |
| <b>Formulation</b>    | Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.<br>Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.<br>Please refer to the specific buffer information in the printed manual. |
| <b>Reconstitution</b> | Please refer to the printed manual for detailed information.   |

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

NKG2-D type II integral membrane protein (NKG2D) is a type II transmembrane glycoprotein which belongs to the CD94/NKG2 family. NKG2D is expressed on natural killer (NK) cells, CD8+ alpha-beta and gamma-delta T-cells. As an activating and costimulatory receptor, it involved in immunosurveillance upon binding to various cellular stress-inducible ligands displayed at the surface of autologous tumor cells and virus-infected cells. It provides both stimulatory and costimulatory innate immune responses on activated killer (NK) cells, leading to cytotoxic activity. It stimulates perforin-mediated elimination of ligand-expressing tumor cells. Signaling involves calcium influx, culminating in the expression of TNF-alpha. NKG2D participates in NK cell-mediated bone marrow graft rejection and survival of NK cells. It Binds to ligands belonging to various subfamilies of MHC class I-related glycoproteins including MICA, MICB, RAET1E, RAET1G, ULBP1, ULBP2, ULBP3 (ULBP2>ULBP1>ULBP3) and ULBP4.

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017