A Reliable Research Partner in Life Science and Medicine

# Recombinant Mouse IL1R1/CD121a Protein (Fc Tag)

Catalog Number: PKSM041047

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Description**

**Species** Mouse

Source HEK293 Cells-derived Mouse IL1R1/CD121a protein Leu20-Lys338, with an C-

terminal Fc

Calculated MW 64.0 kDa Observed MW 85-110 kDa Accession P13504

**Bio-activity** Not validated for activity

### **Properties**

> 95 % as determined by reducing SDS-PAGE. **Purity** 

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

This product is provided as lyophilized powder which is shipped with ice packs. Shipping

Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. **Formulation** 


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

#### Data



> 95 % as determined by reducing SDS-PAGE.

## Background

#### Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

**Elabscience®** 

Mouse Interleukin-1 receptor type 1/IL-1 RI is a cytokine receptor that belongs to the interleukin-1 receptor family. This protein is a receptor for interleukin 1 alpha (IL1A), interleukin 1 beta (IL1B), and interleukin 1 receptor antagonist (IL1R A). It is an important mediator involved in many cytokine induced immune and inflammatory responses. An IL1 receptor accessory protein that can heterodimerize with the Type I receptor in the presence of IL1 $\alpha$  or IL1 $\beta$ but not IL1ra, was identified. This Type I receptor complex appears to mediate all the known IL1 biological responses. The receptor Type II has a short cytoplasmic domain and does not transduce IL1 signals. In addition to the membranebound form of IL1 RII, a naturally occurring soluble form of IL1 RII has been described. It has been suggested that the Type II receptor, either as the membranebound or as the soluble form, serves as a decoy for IL1 and inhibits IL1 action by blocking the binding of IL1 to the signaling Type I receptor complex.

Fax: 1-832-243-6017