# SNRNP25 Polyclonal Antibody

**catalog number:** **E-AB-52686**

**Note:** *Centrifuge before opening to ensure complete recovery of vial contents.*

## Description

| | |
|---|---|
| **Reactivity** | Human;Mouse |
| **Immunogen** | Fusion protein of human SNRNP25 |
| **Host** | Rabbit |
| **Isotype** | Rabbit IgG |
| **Purification** | Antigen affinity purification |
| **Buffer** | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |

| Applications | Recommended Dilution |
|---|---|
| **IHC** | 1:50-1:300 |

## Data



Immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using SNRNP25 Polyclonal Antibody at dilution of 1:55(×200)



Immunohistochemistry of paraffin-embedded Human gastric cancer tissue using SNRNP25 Polyclonal Antibody at dilution of 1:55(×200)

## Preparation & Storage

| | |
|---|---|
| **Storage** | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| **Shipping** | The product is shipped with ice pack,upon receipt,store it immediately at the temperature recommended. |

## Background

Two types of spliceosomes catalyze splicing of pre-mRNAs. The major U2-type spliceosome is found in all eukaryotes and removes U2-type introns, which represent more than 99% of pre-mRNA introns. The minor U12-type spliceosome is found in some eukaryotes and removes U12-type introns, which are rare and have distinct splice consensus signals. The U12-type spliceosome consists of several small nuclear RNAs and associated proteins. This gene encodes a 25K protein that is a component of the U12-type spliceosome.


Toll-free: 1-888-852-8623
Tel: 1-832-243-6086
Fax: 1-832-243-6017
Web: www.elabscience.com
Email: techsupport@elabscience.com