## **Elabscience**®

## Recombinant Mouse IL-19 protein(His Tag)

## Catalog Number: PKSM041507

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Species	Mouse
Source	E.coli-derived Mouse IL-19 protein Leu 25-Ala 176, with an C-terminal His
Calculated MW	18.5 kDa
Observed MW	17-25 kDa
Accession	Q8CJ70
Bio-activity	Measure by its ability to induce proliferation in BaF3 cells transfected with human IL-
	20 R alpha and human IL-20 R beta. The $ED_{50}$ for this effect is <0.6 ng/mL.
Properties	
Purity	> 98 % as determined by reducing SDS-PAGE.
Endotoxin	< 0.1 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of
	reconstituted samples are stable at $<$ -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4.
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants
	before lyophilization.
	Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.
Data	

kDa	10000
75- 63- 48-	
35-	
25-	
17- 11-	

> 98 % as determined by reducing SDS-PAGE.

Background

## **Elabscience**®

The molecular features at the IL19 locus may modestly alter the establishment of HIV-1 infection. Interleukin (IL) 19, IL-20, and IL-24 belong to the IL-10 cytokine family and have been identified to play a role in the regulation of epidermal functions and inflammation. The expression of IL19 in biopsies of patients with active ulcerative colitis was increased compared with patients with quiescent ulcerative colitis and that colitis was attenuated in IL-19-deficient mice. The disruption of the epithelial barrier with dextran sodium sulfate leads to increased IL-19 expression. Attenuated colitis in IL-19-deficient animals was associated with reduced numbers of IL-6-producing macrophages in the inflamed colonic lamina propria. Microbial-driven expression of IL-19 by intestinal macrophages may contribute to the pathogenesis of inflammatory bowel disease.