


Recombinant Mouse PSMA / FOLH1 protein (His Tag)

Catalog Number: PDEM100301

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Species	Mouse
Source	E.coli-derived Mouse PSMA protein Lys45-Phe335, with an N-terminal His
Mol_Mass	31.9 kDa
Accession	O35409
Bio-activity	Not validated for activity
Properties	
Purity	> 95% as determined by reducing SDS-PAGE.
Endotoxin	< 10 EU/mg of the protein as determined by the LAL method
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

Data	
<div> <div> <div>KDa</div> <div>M</div> <div>R</div> </div> <div> <div>80</div> <div>60</div> <div>40</div> <div>30</div> <div>20</div> </div> <div>  </div> </div>	

> 95 % as determined by reducing SDS-PAGE.

Background

Glutamate carboxypeptidase 2, also known as Glutamate carboxypeptidase II, Membrane glutamate carboxypeptidase, Prostate-specific membrane antigen, GCPII, PSMA, FOLH1, and NAALAD1, is a single-pass type I membrane protein which belongs to the peptidase M28 family and M28B subfamily. FOLH1 is highly expressed in prostate epithelium. It is detected in urinary bladder, kidney, testis, ovary, fallopian tube, breast, adrenal gland, liver, esophagus, stomach, small intestine, colon, brain (at protein level), and the capillary endothelium of a variety of tumors. FOLH1 has both folate hydrolase and N-acetylated alpha linked acidic dipeptidase (NAALADase) activity. It has a preference for tri-alpha-glutamate peptides. Genetic variation in FOLH1 may be associated with low folate levels and consequent hyperhomocysteinemia. This condition can result in increased risk of cardiovascular disease, neural tube defects, and cognitive deficits. FOLH1 also shows a promising role in directed imaging and therapy of recurrent or metastatic disease.

For Research Use Only