

Recombinant Human IL-29 protein(His Tag)

Catalog Number: PKSH034114

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

| | |
|----------------------|--|
| Species | Human |
| Source | E.coli-derived Human IL-29 protein Pro 23-Thr 200, with an C-terminal His |
| Calculated MW | 20.7 kDa |
| Observed MW | 22 kDa |
| Accession | Q8IU54 |
| Bio-activity | Measure by its ability to induce IL-8 secretion in HuH7 cells. The ED ₅₀ for this effect is <6 ng/mL. |

Properties

| | |
|-----------------------|--|
| Purity | > 98 % as determined by reducing SDS-PAGE. |
| Endotoxin | < 0.1 EU per µg of the protein as determined by the LAL method. |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. |
| Formulation | Lyophilized from sterile PBS, pH 8.0. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual. |
| Reconstitution | Please refer to the printed manual for detailed information. |

Background

Cytokine with antiviral, antitumour and immunomodulatory activities. Plays a critical role in the antiviral host defense, predominantly in the epithelial tissues. Acts as a ligand for the heterodimeric class II cytokine receptor composed of IL10RB and IFNLR1, and receptor engagement leads to the activation of the JAK/STAT signaling pathway resulting in the expression of IFN-stimulated genes (ISG), which mediate the antiviral state. Has a restricted receptor distribution and therefore restricted targets: is primarily active in epithelial cells and this cell type-selective action is because of the epithelial cell-specific expression of its receptor IFNLR1. Exerts an immunomodulatory effect by up-regulating MHC class I antigen expression.