Recombinant Mouse Angptl4 Protein(Fc Tag)

Catalog Number: PDMM100118

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description		
Species	Mouse	
Source	Mammalian-derived Mouse Angptl4 protein Arg168-Ser410, with an C-terminal Fc	
Calculated MW	49.4 kDa	
Observed MW	60 kDa	
Accession	Q9Z1P8-1	
Bio-activity	Not validated for activity	
Properties		
Purity	> 90% as determined by reducing SDS-PAGE.	
Endotoxin	< 1.0 EU/mg of the protein as determined by the LAL method	
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80	
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20 °C for 3 months.	
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.	
Formulation	Lyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5%	
	Mannitol.	
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of	
	0.5 mg/mL. Concentration is measured by UV-Vis.	

Data

kDa	MR	
80 60	-	
40	-	
30	-	
20	-	

SDS-PAGE analysis of Mouse Angptl4 proteins, 2 μg/lane of Recombinant Mouse Angptl4 proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 49.4 KD

Background

Elabscience®

ANGPTL4, also known as ANGPTL2, is a protein with hypoxia-induced expression in endothelial cells. It contains 1 fibrinogen C-terminal domain and is expressed at high levels in the placenta, heart, liver, muscle, pancreas and lung but expressed poorly in the brain and kidney. ANGPTL4 inhibits proliferation, migration, and tubule formation of endothelial cells and reduces vascular leakage. It may act as a regulator of angiogenesis and modulate tumorigenesis. It inhibits proliferation, migration, and tubule formation of endothelial cells and reduces vascular leakage. It may also exert a protective function on endothelial cells through an endocrine action. ANGPTL4 is directly involved in regulating glucose homeostasis, lipid metabolism, and insulin sensitivity. In response to hypoxia, the unprocessed form of the protein accumulates in the subendothelial extracellular matrix (ECM). The matrix-associated and immobilized unprocessed form limits the formation of actin stress fibers and focal contacts in the adhering endothelial cells and inhibits their adhesion. It also decreases motility of endothelial cells and inhibits the sprouting and tube formation.