



A Reliable Research Partner in Life Science and Medicine

# **Human MMP-9 Antibody Pair Set**

Catalog No. E-KAB-0056 Applications ELISA

**Synonyms** MMP9, CLG4B, Gelatinase B, GELB, MANDP2, 92kDa Type IV Collagenase, 92 KDa

Gelatinase

## Kit components & Storage

Title	Specifications	Storage
Human MMP-9 Capture Antibody	1 vial, 100 μ g	Store at -20°C for one year.
		Avoid freeze / thaw cycles.
Human MMP-9 Detection Antibody	1 vial, 50 μL	Store at -20°C for one year.
(Biotin)		Avoid freeze / thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### **Product Information**

Items		Characteristic (E-KAB-0056)		
		Human MMP-9 Capture Antibody	Human MMP-9 Detection Antibody	
			(Biotin)	
Immunogen	Immunogen	Recombinant Human MMP-9 protein	Recombinant Human MMP-9 protein	
Information Swissprot		P14780		
Product details	Reactivity	Human	Human	
	Host	Mouse	Goat	
	Conjugation	Unconjugated	Biotin	
	Concentration	0.5mg/mL	/	
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300, 1%	
		glycerol, pH 7.4	protective protein, 50% glycerol, pH	
			7.4	
	Purify	Protein A or G	Antigen Affinity	
	Specificity	Detects Human MMP-9 in ELISAs.		

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: <a href="mailto:www.elabscience.com">www.elabscience.com</a> Email: <a href="mailto:techsupport@elabscience.com">techsupport@elabscience.com</a>





A Reliable Research Partner in Life Science and Medicine

## **Applications**

Human MMP-9 Sandwich ELISA Assay:

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4μg/mL	Human MMP-9 Capture Antibody	
Capture			Alisa
ELISA	1:1000-1:10000	Human MMP-9 Detection	Optical Density
Detection		Antibody (Biotin)	0.01 0.01 1 10 100 Human MMP-9 concentration(ng/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

### **Background**

Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. The enzyme encoded by this gene degrades type IV and V collagens. Studies in rhesus monkeys suggest that the enzyme is involved in IL-8-induced mobilization of hematopoietic progenitor cells from bone marrow, and murine studies suggest a role in tumorassociated tissue remodeling.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: <a href="mailto:www.elabscience.com">www.elabscience.com</a> Email: <a href="mailto:techsupport@elabscience.com">techsupport@elabscience.com</a>